

Proliferation (CFU/ml) X10 ⁵	+ bacteria	+bacteria + BC	+bacteria +egg	+bacteria +BC+egg
<i>E.coli</i> K12	54.3 +/- 1.3	50.7 +/- 2.4	54.0 +/- 3.2	50.0 +/- 2.5
<i>E.coli</i>	55.3 +/- 0.8	50.2 +/- 2.1	56.5 +/- 3.3	50.3 +/- 2.2
EPEC	56.5 +/- 2.5	57.2 +/- 2.8	57.5 +/- 1.3	57.7 +/- 2.6
<i>Salmonella</i>	59.5 +/- 2.3	61.5 +/- 1.8	61.3 +/- 3.2	60.3 +/- 1.9
<i>Klebsiella</i>	59.3 +/- 0.8	61.7 +/- 2.2	61.5 +/- 2.5	61.7 +/- 1.1
<i>Enterococcus</i>	62.7 +/- 2.0	63.5 +/- 0.6	63.7 +/- 0.9	62.7 +/- 0.6
<i>Proteus</i>	51.7 +/- 1.9	52.3 +/- 1.2	52.3 +/- 0.6	51.5 +/- 0.8
<i>Staphylococcus</i>	64.2 +/- 0.8	61.3 +/- 2.4	65.3 +/- 2.2	62.0 +/- 2.6
<i>Streptococcus</i>	43.2 +/- 1.5	40.8 +/- 1.4	44.5 +/- 0.8	40.2 +/- 1.2

Supplemental Table 1. (Lack of) effect of Bovine Colostrum (BC), egg or BC+egg combination on bacterial growth.

Data shown for 24h timepoint using 5 mg/ml of BC alone, 5 mg/ml egg alone or same final concentrate (5 mg/ml) of a 60:40 BC:egg mixture. Results expressed as mean +/- SEM of 6 wells.