

Suppressive Activity of *Glechoma hederacea* Extracts against the Phytopathogenic Oomycete *Plasmopara viticola*, and First Screening of the Active Metabolites

Jesús G. Zorrilla ^{1,2}, Oscar Giovannini ³, Stefano Nadalini ^{3,4}, Alberto Zanini ⁴, Maria Teresa Russo ¹, Marco Masi ^{1*}, Gerardo Puopolo ^{3,4*}, and Alessio Cimmino ¹

¹ Department of Chemical Sciences, University of Naples Federico II, Complesso Universitario Monte S. Angelo, Via Cintia, 80126 Naples, Italy; mariateresa.russo2@unina.it (M.T.R.); alessio.cimmino@unina.it (A.C.)

² Allelopathy Group, Department of Organic Chemistry, Facultad de Ciencias, Institute of Biomolecules (INBIO), University of Cadiz, C/Avenida República Saharaui, s/n, 11510 Puerto Real, Spain; jesus.zorrilla@uca.es (J.G.Z.)

³ Research and Innovation Centre, Fondazione Edmund Mach, Via E. Mach 1, San Michele all'Adige 38098, Italy; oscar.giovannini@fmach.it (O.G.)

⁴ Center Agriculture Food Environment (C3A), University of Trento, Via E. Mach 1, San Michele all'Adige 38098, Italy; stefano.nadalini@unitn.it (S.N.); alberto.zanini@studenti.unitn.it (A.Z.)

* Correspondence: marco.masi@unina.it (M.M.); gerardo.puopolo@unitn.it (G.P.)

Index

Figure S1. ¹ H NMR spectrum of carvacrol in CDCl ₃ (400 MHz)	2
Figure S2. ¹ H NMR spectrum of cirsimarinin in CD ₃ OD (400 MHz).....	2
Figure S3. ¹ H NMR spectrum of apigenin in CD ₃ OD (400 MHz)	3
Figure S4. ¹ H NMR spectrum of methyl caffeate in CD ₃ OD (400 MHz)	3
Figure S5. ¹ H NMR spectrum of caffeic acid in CD ₃ OD (400 MHz)	4
Figure S6. ¹ H NMR spectrum of rosmarinic acid in CD ₃ OD (400 MHz).....	4
Figure S7. Control of <i>Plasmopara viticola</i> on grapevine leaf discs through the application of <i>Glechoma hederacea</i> aqueous extracts.....	5




Figure S1. ^1H NMR spectrum of carvacrol in CDCl_3 (400 MHz)




Figure S2. ^1H NMR spectrum of cirsimarinin in CD_3OD (400 MHz)




Figure S3. ^1H NMR spectrum of apigenin in CD_3OD (400 MHz)




Figure S4. ^1H NMR spectrum of methyl caffeoate in CD_3OD (400 MHz)




Figure S5. ^1H NMR spectrum of caffeic acid in CD_3OD (400 MHz)




Figure S6. ^1H NMR spectrum of rosmarinic acid in CD_3OD (400 MHz)




Figure S7. Control of *Plasmopara viticola* on grapevine leaf discs through the application of *Glechoma hederacea* aqueous extracts. Representative photographs showing the effect of the application of treatments in controlling *P. viticola* on grapevine leaf discs. The treatments evaluated were: (1) grapevine leaf discs treated with distilled water; (2) grapevine leaf discs treated with copper hydroxide (2 g/L) and (3) grapevine leaf discs treated with *G. hederacea* aqueous extracts.