



Proceeding Paper

Quality of Surface and Ground Water in Three States of Nigeria: Assessment of Physicochemical Characteristics and Selected Contamination Patterns [†]

Francis Olawale Abulude ^{1,*}, Akinyinka Akinnusotu ² , Ebenezer Alaba Adeoya ³, Samson Olatunde Mabayoje ⁴, Samuel Dare Oluwagbayide ⁵, Kikelomo Mabinuola Arifalo ⁶ and Ademola Adamu ⁶

¹ Environmental and Sustainable Research Group, Science and Education Development Institute, Akure 340001, Ondo State, Nigeria

² Department of Science Laboratory Technology, Rufus Giwa Polytechnic, Owo 341101, Ondo State, Nigeria; akinnusotuakinyinka@gmail.com

³ National Agency for Science and Engineering Infrastructure (NASENI), Akure 340001, Ondo State, Nigeria; adeoyaeben@yahoo.com

⁴ Department of Biological Sciences, College of Sciences, Afe Babalola University (ABUAD), Ado Ekiti 360101, Ekiti State, Nigeria; mabayojeso@abuad.edu.ng

⁵ Department of Agricultural and Bio-Environmental Engineering, Federal Polytechnic, P.M.B 50, Ilaro 112106, Ogun State, Nigeria; samuel.oluwagbayide@federalpolyilaro.edu.ng

⁶ Department of Chemistry, Bamidele Olomilua University of Education, Science and Technology, Ikere Ekiti 361101, Ekiti State, Nigeria; karifalo@yahoo.co.uk (K.M.A.); adamu.ademola@coeikere.edu.ng (A.A.)

* Correspondence: waleabul@outlook.com; Tel.: +234-803445674

† Presented at the 7th International Electronic Conference on Water Sciences, 15–30 March 2023; Available online: <https://ecws-7.sciforum.net>.



Citation: Abulude, F.O.; Akinnusotu, A.; Adeoya, E.A.; Mabayoje, S.O.; Oluwagbayide, S.D.; Arifalo, K.M.; Adamu, A. Quality of Surface and Ground Water in Three States of Nigeria: Assessment of Physicochemical Characteristics and Selected Contamination Patterns. *Environ. Sci. Proc.* **2023**, *25*, 48. <https://doi.org/10.3390/ECWS-7-14258>

Academic Editor: Athanasios Loukas

Published: 16 March 2023



Copyright: © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Abstract: The study of water quality is crucial given the amount of industrial, agricultural, and other human activities at the sampling sites. The aim of the study was to assess the physicochemical characteristics and selected contamination patterns of water samples in Nigeria. This study used conventional analytical techniques to analyze the physicochemical parameters in water samples from 33 sampling sites (dug wells, boreholes, rivers, and rainwater) in three different states (Ekiti, Osun, and Ondo) of Nigeria. These parameters included pH, total dissolved solids (TDS), electrical conductivity (EC), temperature, relative humidity (RH), and four elemental parameters (Ca, Na, Fe, and Cu). The enrichment factor (EF), contamination factor (CF), and metal index (MI) were used to characterize the data. Temperature (28.17 °C), TDS (130.2 mg/L), EC (260.0 μS/cm), pH (6.88), Na (14.47 ppm), Ca (25.74 ppm), Fe (0.49 ppm), and Cu (0.08 ppm) were the average values from the results. Na and Ca had a direct relationship with one another. The levels of heavy metals were below those recommended by the Nigerian Industrial Standard for Drinking Water Quality (NISDQW) and the World Health Organization (WHO). The metal levels in the water samples were over 1.5, which is the threshold value indicated by the EF classifications. In particular, EFs were moderate to significantly enriched. All element CFs were below the Level 1 pollution threshold. The water samples are pure based on the MI's rating of water quality. Human and natural activities may represent a risk to the local public health; hence, it is highly advisable that all stakeholders adopt rapid and long-lasting collective action to limit pollution levels as part of the water quality governance system.

Keywords: anthropogenic activities; water samples; heavy metals; TDS; enrichment factor; WHO; SON

1. Introduction

The accessibility and sustainability of water and sanitation for every person worldwide by 2030 are among the key sustainable development objectives (Goal 6) [1,2]. The

objectives are to provide everyone with equitable access to clean, inexpensive drinking water; achieve adequate and equitable access to sanitation and hygiene; improve water quality by reducing pollution, eliminating dumping, and minimizing the release of hazardous chemicals and materials; achieve half as much untreated wastewater; implement integrated water resources management at all levels; and achieve protection and restoration [3].

For continued population expansion and development, access to a secure and reliable water supply is a crucial requirement [4]. Both surface and ground water are essential sources of water for the global population. About 90% of the world's readily usable freshwater resources are found in groundwater, with the other 10% being found in lakes, reservoirs, rivers, and wetlands. Moreover, the expansion of an estimated 40% of the world's agricultural production is supported by groundwater irrigation of arable lands [4]. Groundwater is the most dependable source of drinking water in sub-Saharan Africa [5].

Southwest Nigeria faces a number of difficulties, including how to accomplish sustainable development goals and provide drinkable water for its expanding population. Due to the scarcity of surface water supplies in some regions, people are forced to use underground water, which presents a problem in those areas. In the states we examined, groundwater and surface water are significant natural resources that have an impact on both human and animal health and welfare. As a result, primary research and quality control efforts should be directed at the quality of these resources. This premise would not be considered out of place if the qualities of the water samples are determined. The main goal of this study was to assess the quality (physicochemical characteristics and selected contamination patterns) of ground and surface water samples (borehole, dug well, rainwater, and river) from various communities in chosen areas of three states (Osun, Ondo, and Ekiti) in the southwest of Nigeria.

2. Materials and Methods

The Osun, Ekiti, and Ondo States of Nigeria, which make up the study area, are situated in the southwest of the country. The research area's climate generally follows the same pattern. In terms of population, transportation, industry, housing, and agriculture, the states are rapidly expanding. For this experiment in November 2022, 33 water samples were selected from rivers (10), hand-dug wells (14), boreholes (7), and rain (2). The samples were obtained in polyethylene bottles, cleaned properly with distilled water, and then treated with nitric acid before being filtered through membrane filters with pores measuring 0.45 microns. A Garmin global positioning system was used to determine the coordinates of the sampling locations. A portable multi-parameter meter called the Temp/pH/TDS/EC meter (model MI 1399) was used to measure the temperature, electrical conductivity, pH, temperature, and TDS in situ immediately after sample collection in the field. To stop the precipitation of trace elements, nitric acid was used to acidify the water samples. The elements (Na, Ca, Cu, and Fe) were evaluated utilizing conventional methods (AAS, Buck Scientific GVP 210, USA) in the Central Laboratory of Quality Monitoring at Afe Babalola University in Ado-Ekiti, Ekiti State, Nigeria. The metal index, contamination factor, and enrichment factor (EF) were determined. EF was calculated using this factor [6]:

$$EF = \frac{(C_i/C_{ref}) \text{ sample}}{(B_i/B_{ref}) \text{ Background}} \quad (1)$$

where B_i is the background value of an element of interest and B_{ref} is the background value of the reference element in the study area, C_i is the concentration of trace elements in the sample, C_{ref} is the concentration of the reference element in the sample, and C_{ref} is the background value of the reference analyte in the sample. The reference element used in this study was Fe, which is most widely used for normalization [6]. EF classification: $EF < 2$ (deficiency to minimal enrichment), $2 \leq EF < 5$ (moderate enrichment), $5 \leq EF < 20$ (significant enrichment), $20 \leq EF < 40$ (very high enrichment), and $EF \geq 40$ (extremely high enrichment).

CF was calculated using this factor [6]:

$$CF = C_i/B_i \tag{2}$$

These values were obtained by calculating the ratio of the element’s background concentration to the concentration of the element present in the sample [7]. C_i = concentration of the examined element i , and B_i = geochemical background value of the element. The contamination values in increasing order of contaminations are 0 = none, 1 = none to medium, 2 = moderate, 3 = moderate to strong, 4 = strongly polluted, 5 = strong to very strong, 6 = very strong [8]. In terms of metal and metalloid contamination, MI shows an overall trend in water quality [9], where H_c is the i th parameter’s monitored value (in mg/L), and H_{mac} is the i th parameter’s maximum permissible concentration [10]. According to the MI, the water is either lowly polluted ($MI < 10$) or moderately polluted ($10 < MI < 20$) [11]. MI was calculated by Equation (3) [11]:

$$MI = \sum_{i=1}^n \frac{H_c}{H_{mac}} \tag{3}$$

The MI classification for water samples are: <0.3 = very pure (Class I), $0.3-1.0$ pure (Class II), $1.0-2.0$ slightly affected (Class III), $2.0-4.0$ moderately affected (Class IV), $4.0-6.0$ strongly affected (Class V), >6.0 seriously affected (Class VI).

3. Results and Discussion

The recorded mean pH levels of the water samples did not differ from one another statistically ($p > 0.05$). The pH in the water samples ranged from 5.66 to 7.89, with a mean value of 6.88 0.60 (Table 1). Dug wells had the lowest pH readings, whereas boreholes had the highest. Except for the lowest pH level, the pH of water was within the range of 6.5–8.5 allowed by WHO [12] and SON [13] for drinking water. The outcome is consistent with the findings of Appiah-Opong et al. [14]. Water with a pH level below 6.5 is considered to be too acidic for human consumption, which could lead to conditions such as acidosis and harm the digestive and lymphatic systems [6]. Statistical analysis revealed that the means were not different ($p > 0.05$) from one another, despite the fact that the electrical conductivity of the water was generally higher in the dug wells than in other samples. The mean EC values in the water samples ranged thus: dug well (84–1003 μ S/cm), borehole (136–386 μ S/cm), rain (54–59 μ S/cm), and rivers (76–297 μ S/cm). The overall mean was 260 μ S/cm.

Table 1. Basic Description of the Parameters.

Variable	Mean	SE Mean	StDev	CoefVar (%)	Min	Q1	Median	Q3	Max	Skewness	Kurtosis
Temperature (°C)	28.17	0.28	1.67	5.93	24.80	26.90	28.60	29.30	31.20	−0.38	−0.72
TDS (mg/L)	130.20	15.60	89.80	68.99	27.00	71.00	118.00	162.00	511.00	2.49	9.39
EC (μ S/cm)	260.00	30.70	176.40	67.86	54.00	144.00	239.00	314.50	1003.00	2.45	9.10
pH	6.88	0.10	0.60	8.00	5.66	6.47	7.02	7.27	7.89	−0.52	−0.46
Na (mg/L)	14.47	0.63	3.62	24.98	9.40	11.05	14.80	17.40	23.50	0.46	−0.36
Ca (mg/L)	25.74	1.74	10.02	38.92	8.20	17.80	22.90	31.15	48.30	0.69	−0.18
Fe (mg/L)	0.49	0.05	0.30	60.14	0.18	0.29	0.40	0.62	1.35	1.62	2.29
Cu (mg/L)	0.08	0.01	0.04	41.48	0.02	0.06	0.08	0.11	0.16	0.57	−0.27

WHO (2011)—temperature (22–29 °C), TDS (1000 mg/L), EC 1000 (mg/L), pH (6.5–8.5), Na (200), Ca (75), Fe (0.03), Cu (200 mg/L); NISDQW (2015)—temperature (22–29 °C), TDS (1000 mg/L), EC 1000 (mg/L), pH (6.5–8.5), Na (200), Ca (75 mg/L), Fe (0.03), Cu (200 mg/L).

The EC results fell below the 1000 μ S/cm drinking water limit set by WHO [12] and SON [13]. Although EC is not a concern for human or aquatic health, it might be a sign of other issues with water quality [14]. The high values of EC in the dug well could be linked to anthropogenic activities, as well as the soil’s mineral or salt dissolution [15]. TDS values ranged between 27 and 511 mg/L, with 260 mg/L being the mean. The considerable high variability in the water samples was shown by the coefficient of variation

in terms of percentages. No differences in water temperature were found to be statistically significant ($p > 0.05$). The results showed the range, standard error, skewness, and kurtosis of temperature as 24.80–31.20 °C, 0.28, −0.38, and −0.72, respectively. The minimum value was recorded in a stream. The reason could be due to the activities of the sampling which assisted in the aeration of the water. In addition to the time of the sample, other factors that may affect temperature include water depth, season, groundwater influx, and air circulation [16]. These findings concur with those made in the Ivory Coast by Koffi et al. [15].

The concentrations of Na, Ca, Fe, and Cu in the thirty-three sampling points ranged from 9.40 to 23.50 mg/L, 8.20 to 48.30 mg/L, 0.18 to 1.35 mg/L, and 0.02 to 0.16 mg/L, respectively. The elements' average concentrations fell in the following order: Ca > Na > Fe > Cu (Table 1). These results agree with those found by Koffi et al. [15] in the Ivory Coast. The number of components was below what is considered to be acceptable for drinking and irrigation water on a national and international level. Only a few of the water samples had iron contents that were above the WHO limit (0.3 mg/L); over 90% of the samples were regarded as suitable for use in irrigation and human consumption. Undesirable tastes and smells are typically connected to underground water that contains more iron [17]. It is possible that the iron in water samples came from natural sources as well [18]. The concentration of copper was higher in river water samples than in others. This shows that the rivers are either naturally high in copper or absorbed from soils with Cu fertilizers. The calcium concentration was greater than the amount of sodium content for more than 90% of the water samples gathered. The amount of carbonate minerals that make up the water-bearing formations, ion exchange mechanisms, and the precipitation of calcite in the aquifer can all be used to explain this [19].

Enrichment factors of elements followed this order Ca > Cu > Na > Fe. The metal levels in the water samples were over 1.5, which is the threshold value indicated by the EF classifications. In particular, EFs were moderately to significantly enriched. Figure 1 displays the water sample histogram. The samples fell within the very pure, Class I category. This result was compared with the findings of Khosnam et al. [20] for the Silakhor River in Iran.




Figure 1. Cont.




Figure 1. The Metal Index, Contamination Factors, and Enrichment Factor of the Water Samples.

Figure 2 and Table 2 depicted the matrix correlation (Pearson correlation). This showed the correlations between the physicochemical parameters and the elements. There were strong positive relationships between EC and TDS ($r = 0.99$) and Ca and Na ($r = 0.72$) depicting that an increase in EC causes an increase in TDS, showing a direct relationship between the variables. Ca and Na had a substantial correlation, which suggested that the components in the water samples might have had identical origins.




Figure 2. The matrix correlation of the water samples (Pearson correlation).

Table 2. The correlation of the water samples (Pearson correlation).

EC	0.99						
pH	−0.00	−0.01					
Na	0.12	0.12	0.15				
Ca	0.01	0.01	0.10	0.72			
Fe	−0.30	−0.29	−0.19	0.20	0.20		
Cu	0.22	0.22	−0.13	−0.07	−0.09	−0.03	
	TDS	EC	pH	Na	Ca	Fe	

4. Conclusions

This study assessed the physicochemical characteristics and selected contamination patterns of thirty-three water samples from three southwest states in Nigeria. Additionally, in the study, the EF, CF, and MI were determined. The physicochemical and elemental levels obtained were within the standard limits of WHO and NISDQW. The EFs of the water samples were moderately to significantly enriched. All elemental CFs were below the Level 1 pollution threshold. The MI rating showed that they are suitable for human and animal consumption and also for irrigation purposes. Although the present study concluded that the water samples were good, efforts should be put in place to prevent them from being polluted.

Author Contributions: Conceptualization, F.O.A.; methodology, A.A. (Akinyinka Akinnusotu) and S.D.O.; formal analysis, E.A.A. and S.D.O.; investigation, S.O.M.; data curation, K.M.A.; writing—original draft preparation, F.O.A.; writing—review and editing, E.A.A., K.M.A., S.D.O. and A.A. (Akinyinka Akinnusotu); supervision, A.A. (Ademola Adamu) and S.O.M. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: The raw data are available upon request. Please contact the corresponding authors.

Acknowledgments: The authors are grateful for the permissions granted by the owners of the dug wells and boreholes to obtain samples from their homes and offices.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Abulude, F.O.; Fagbayide, S.D. Water, sanitation, and poverty in the changing world. Case of Nigeria. *An. Univ. Din Oradea Ser. Geogr.* **2018**, *28*, 91–96.
2. Oluwagbayide, S.D.; Abulude, F.O. Assessment of the quality of dug well water samples in Nigeria and their suitability for drinking and irrigation purposes. *Sustain. Water Resour. Manag.* **2022**, *8*, 149. [CrossRef]
3. United Nations. Department of Economic and Social Affairs Disability. Envision2030 Goal 6: Clean Water and Sanitation. Targets. Available online: <https://www.un.org/development/desa/disabilities/envision2030-goal6.html> (accessed on 24 November 2020).
4. Saana, S.B.B.M.; Fosu, S.A.; Sebiawu, G.E.; Jackson, N.; Karikari, T. Assessment of the quality of groundwater for drinking purposes in the Upper West and Northern regions of Ghana. *SpringerPlus* **2016**, *5*, 2001. [CrossRef] [PubMed]
5. Iyasele, J.U.; Idiata, D.J. Physico-chemical and microbial analysis of borehole water samples: A case of some boreholes in Edo North, Edo state. *J. Emerg. Trends Eng. Appl. Sci.* **2011**, *2*, 1064–1067.
6. Bhuiyan, M.A.H.; Parvez, L.; Islam, M.A.; Dampare, S.B.; Suzuki, S. Heavy metal pollution of coal mine-affected agricultural soils in the northern part of Bangladesh. *J. Hazard. Mater.* **2010**, *173*, 384–392. [CrossRef] [PubMed]
7. Hakanson, L. An ecological risk index for aquatic pollution control. A sediment logical approach. *Water Res.* **1980**, *14*, 975–1001. [CrossRef]
8. Varol, M. Assessment of heavy metal contamination in sediments of the Tigris River (Turkey) using pollution indices and multivariate statistical techniques. *J. Hazard. Mater.* **2011**, *195*, 355–364. [CrossRef] [PubMed]
9. Edet, A.E.; Offiong, O.E. Evaluation of water quality pollution indices for heavy metal contamination monitoring. A study case from Akpabuyo-Odukpani area, Lower Cross River Basin (southeastern Nigeria). *GeoJournal* **2002**, *57*, 295–304. [CrossRef]
10. Shil, S.; Singh, U.K. Health risk assessment and spatial variations of dissolved heavy metals and metalloids in a tropical river basin system. *Ecol. Indic.* **2019**, *106*, 105455. [CrossRef]
11. Rezaei, A.; Hassani, H.; Hassani, S.; Jabbari, N.; Fard Mousavi, S.B.; Rezaei, S. Evaluation of groundwater quality and heavy metal pollution indices in Bazman basin, southeastern Iran. *Groundw. Sustain. Dev.* **2019**, *9*, 100245. [CrossRef]
12. WHO. *Guidelines for Drinking-Water Quality*, 4th ed.; WHO: Rome, Italy, 2011. Available online: http://www.who.int/about/licensing/copyright_form/en/index.html (accessed on 12 December 2022).
13. NIS 554: 2015; Nigerian Standard for Drinking Water Quality (NSDWQ). Standard Organization of Nigeria: Abuja, Nigeria, 2015. Available online: <https://africacheck.org/wp-content/uploads/2018/06/Nigerian-Standard-for-Drinking-Water-Quality-NIS-554-2015.pdf> (accessed on 12 December 2022).

14. Appiah-Opong, R.; Ofori, A.; Odosuhene, M.; Ofori-Attach, E.; Nunoo, F.K.E.; Tuffour, I.; Gordo, C.; Arhinful, D.K.; Nyarko, A.K.; Fosu-Mensah, B.Y. Heavy metals concentration and pollution index (HPI) in drinking water along the southwest coast of Ghana. *Appl. Water Sci.* **2021**, *11*, 57. [[CrossRef](#)]
15. Koffi, K.M.; Coulibaly, S.; Atse, B.C.; Paul, E. Survey of heavy metals concentrations in water and sediments of the estuary Bietri Bay. Ebrie Lagoon, Cote D'Ivoire. *Int. J. Res. Earth Environ. Sci.* **2014**, *1*, 2311.
16. Dan, S.F.; Umoh, U.U.; Osabor, V.N. Seasonal variation of enrichment and contamination of heavy metals in the surface water of Qua Iboe River Estuary and adjoining creeks, South-South Nigeria. *J. Oceanogr. Mar. Sci.* **2014**, *5*, 45–54. [[CrossRef](#)]
17. Sethy, S.N.; Syed, T.H.; Kumar, A.; Sinha, D. Hydrogeochemical characterization and quality assessment of groundwater in parts of Southern Gangetic Plain. *Environ. Earth Sci.* **2016**, *75*, 232. [[CrossRef](#)]
18. Smedley, P.L.; Edmunds, W.M.; West, J.M.; Gardner, S.J.; Pelig-Ba, K.B. *Vulnerability of Shallow Groundwater Quality Due to Natural Geochemical Environment. 2, Health Problems Related to Groundwater in the Obuasi and Bolgatanga Areas, Ghana*; British Geological Survey: Nottingham, UK, 1995; p. 130.
19. Sebiawu, E.G.; Fosu, S.A.; Saana, S.B.B.M. A physico-chemical and bacteriological analysis of borehole water samples from the Wa Municipality of the Upper West Region, Ghana. *Int. J. Eng. Res.* **2014**, *3*, 1959–1965.
20. Khoshnam, Z.; Sarikhani, R.; Ghassemi Dehnavi, A.; Ahmadnejad, Z. Evaluation of Water Quality Using Heavy Metal Index and Multivariate Statistical Analysis in Lorestan Province. *Iran. J Adv. Environ. Health Res.* **2017**, *5*, 29–37.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.