

## Occurrence of Picobirnavirus in Domestic and Wild Animals from Three Cities of Brazilian Amazon

**Table S2**

**Table 1.** Data referring to statistical analysis of the frequency of PBV in fecal samples of animals, according to the group, gender, and city of origin, in the period from October-2014 to April-2016

CATEGORIES	PICOBIRNAVIRUS				Total		p
	Positive		Negative		n	%	
	n	%	n	%			
GROUP							0.000 <sup>#</sup>
Avian	2	4.9	39	95.1	41	100.0	
Bovine	3	13.0	20	87.0	23	100.0	
Canine	5	14.7	29	85.3	34	100.0	
Equine	3	7.3	38	92.7	41	100.0	
Feline*	9	27.3	24	72.7	33	100.0	
Marsupial*	1	4.0	24	96	25	100.0	
Bats	0	0.0	30	100.0	30	100.0	
Rodent*	0	0.0	8	100.0	8	100.0	
Swine*	9	39.1	14	60.9	23	100.0	
GENDER							0.918 <sup>¥</sup>
Female	14	14.0	86	86.0	100	100.0	
Male	15	13,5	96	86,5	111	100,0	
CITY							0,004 <sup>¥</sup>
Peixe-Boi*	8	27,6	21	72,4	29	100,0	
Santa Bárbara	10	19,6	41	80,4	51	100,0	
Viseu*	14	7,9	164	92,1	178	100,0	
Total	32	12,4	226	87,6	258	100,0	

p: Probability score of the applied statistic test. <sup>#</sup>G Test. <sup>¥</sup>Chi-square Test. \*Categories that contribute to statistical significance. The p-value was 0.000954.